

Free energy and θ dependence of $SU(N)$ gauge theories*

L. Del Debbio^a, H. Panagopoulos^b, E. Vicari^a

^aDipartimento di Fisica, Università di Pisa, and INFN, I-56127 Pisa, Italy

^bDepartment of Physics, University of Cyprus, Nicosia CY-1678, Cyprus

We study the dependence of the free energy on the CP violating angle θ , in four-dimensional $SU(N)$ gauge theories with $N \geq 3$, and in the large- N limit. Using the Wilson lattice formulation for numerical simulations, we compute the first few terms of the expansion of the ground-state energy $F(\theta)$ around $\theta = 0$, $F(\theta) - F(0) = A_2 \theta^2 (1 + b_2 \theta^2 + \dots)$. Our results support Witten's conjecture: $F(\theta) - F(0) = \mathcal{A} \theta^2 + O(1/N)$ for $\theta < \pi$. We verify that the topological susceptibility has a nonzero large- N limit $\chi_\infty = 2\mathcal{A}$ with corrections of $O(1/N^2)$, in substantial agreement with the Witten-Veneziano formula which relates χ_∞ to the η' mass. Furthermore, higher order terms in θ are suppressed; in particular, the $O(\theta^4)$ term b_2 (related to the $\eta' - \eta'$ elastic scattering amplitude) turns out to be quite small: $b_2 = -0.023(7)$ for $N = 3$, and its absolute value decreases with increasing N , consistently with the expectation $b_2 = O(1/N^2)$.

1. INTRODUCTION

Interest in $SU(N)$ gauge theories has recently been revived by studies in the context of M-theory, AdS/CFT correspondence, and $\mathcal{N} = 2$ SUSY broken to $\mathcal{N} = 1$. A number of predictions have been put forth, and we have set out to check these predictions by numerical simulation.

One aspect of these studies regards the spectrum of confining strings in four-dimensional $SU(N)$ gauge theories. A considerable amount of numerical work has been done on this subject (see [1,2] and references therein). In particular, in Ref. [2] we have presented computations of the string tensions σ_k , related to sources with Z_N charge k , for $SU(4)$ and $SU(6)$ gauge theories. Our results are consistent with the sine formula $\sigma_k/\sigma_1 = \sin(k\pi/N)/\sin(\pi/N)$, with an accuracy of approximately 1% [2%] for $SU(4)$ [$SU(6)$]; this formula emerges in various realizations of supersymmetric $SU(N)$ gauge theories, and in the context of M-theory. On the other hand, our results show deviations from Casimir scaling $\sigma_k/\sigma = k(N-k)/(N-1)$, as might be expected from the short distance behaviour of the potential, or from a strong-coupling limit.

In the present work, we study the θ dependence of $SU(N)$ gauge theories, for $N \geq 3$ and in the large- N limit, by numerical simulation; for a longer write-up of our work, see [3]. The angle θ appears in the Euclidean Lagrangian as:

$$\mathcal{L}_\theta = \frac{1}{4} F_{\mu\nu}^a(x) F_{\mu\nu}^a(x) - i\theta q(x) \quad (1)$$

where $q(x)$ is the topological charge density

$$q(x) = \frac{g^2}{64\pi^2} \epsilon_{\mu\nu\rho\sigma} F_{\mu\nu}^a(x) F_{\rho\sigma}^a(x). \quad (2)$$

The partition function reads:

$$Z(\theta) = \int [dA] \exp(- \int d^4x \mathcal{L}_\theta) \equiv \exp[-V F(\theta)] \quad (3)$$

where $F(\theta)$ is the free (ground state) energy. Witten has argued [4] that in the large- N limit $F(\theta)$ is a multibranched function of the type

$$F(\theta) = N^2 \min_k H[(\theta + 2\pi k)/N] \quad (4)$$

The conjecture was refined [5] and simplified:

$$\Delta F(\theta) = \mathcal{A} \min_k (\theta + 2\pi k)^2 + O(1/N). \quad (5)$$

($\Delta F(\theta) \equiv F(\theta) - F(0)$). In particular, for $\theta < \pi$,

$$\Delta F(\theta) = \mathcal{A} \theta^2 + O(1/N). \quad (6)$$

This conjecture has been supported by gauge/string duality arguments [5].

*Talk presented by H. Panagopoulos

Monte Carlo studies of the θ -dependence are made very difficult by the complex nature of the θ term. In fact $\theta \neq 0$ cannot be directly simulated on the lattice. Here we restrict ourselves to relatively small θ values, where one may expand:

$$\Delta F(\theta) = A_2\theta^2 + A_4\theta^4 + \dots \quad (7)$$

A_2 gives the topological susceptibility, i.e.

$$A_2 = \chi/2, \quad \chi = \langle Q^2 \rangle_{\theta=0}/V, \quad (8)$$

where Q is the topological charge $Q = \int d^4x q(x)$. Higher order coefficients in Eq. (7) can be related to higher moments of the probability distribution $p(Q)$ of the topological charge. For instance,

$$A_4 = -\frac{1}{24V} [\langle Q^4 \rangle - 3\langle Q^2 \rangle^2]_{\theta=0}. \quad (9)$$

A convenient parameterization of the free energy using dimensionless scaling coefficients is:

$$\sigma^{-2} \Delta F(\theta) = \frac{1}{2} C \theta^2 (1 + b_2 \theta^2 + b_4 \theta^4 + \dots) \quad (10)$$

(σ : string tension, $C = \chi/\sigma^2$, $b_2 = A_4/A_2$).

Witten's conjecture implies that C has a finite nonzero large- N limit C_∞ and $b_{2i} = O(1/N^2)$. In particular one expects that b_2 is small, and that it should rapidly decrease with increasing N . A nonzero value of χ_∞ is essential to resolve the $U(1)_A$ problem, and is related to the η' mass [6]:

$$\chi_\infty = 2A = \frac{f_\pi^2 m_{\eta'}^2}{4N_f} + O(1/N). \quad (11)$$

2. CALCULATION AND RESULTS

We present results for four-dimensional $SU(N)$ gauge theories with $N = 3, 4, 6$. In a nutshell, we: Determine the ratio $C = \chi/\sigma^2$; verify the expected behavior $C \sim A + B/N^2$, determining the large- N limit by a fit; obtain a rather accurate estimate of C_∞ ; estimate the values of b_2 , which are found to be very small and rapidly decreasing with N , in support of Witten's conjecture.

Simulation Details: • We use the standard Wilson gauge action and the Cabibbo-Marinari updating algorithm (one $SU(N)$ update $\rightarrow N(N-1)/2$ $SU(2)$ subgroup updates). • Microcanonical

over-relaxation and heat-bath updates are used in a 4:1 ratio. • Values for β are chosen in the weak-coupling, (beyond the 1st order phase transition ($N = 6$), beyond crossover (peak of specific heat $N = 3, 4$)). • Topological quantities are computed via cooling. We measure Q typically every 100 sweeps. • We have employed different lattice sizes, so that finite size effects are under control. Tables of our numerical results can be found in Ref. [3].




FIGURE 1. Autocorrelation time τ_Q versus $\xi_\sigma \equiv \sigma^{-1/2}$. Dotted lines show the linear fits.

A severe form of critical slowing down is seen in the measurement of Q , at large values of N . Our results (Fig. 1) suggest an exponential behavior:

$$\ln \tau_Q \approx N(e_N \xi_\sigma + c_N) \quad (12)$$

($\xi_\sigma \equiv \sigma^{-1/2}$, and e_N, c_N are constants with only weak N -dependence). Thus, $\ln \tau_Q \propto N$ (at fixed ξ_σ). This phenomenon is possibly due to the free-energy barriers present. There are no a priori arguments in favor of this behavior; it simply arises naturally from fits, unlike a power law. No similar effect is observed in plaquette or Polyakov line correlations; this suggests a decoupling between topological modes and nontopological ones, such as those determining confining properties.

In Fig. 2, the ratio C is plotted versus σ , to evidetiate possible scaling corrections (expected: $\mathcal{O}(a^2)$, logs). Corrections are indeed observed for $N = 3$ and $N = 4$, and we must allow for them in a fit, for a continuum extrapolation. Assuming

a linear behavior: $C = C_{\text{cont}} + u\sigma$, to take into account the leading $\mathcal{O}(a^2)$ correction, we obtain:

$$C_{\text{cont}} = 0.0282(12), u = 0.077(24) \quad (N=3) \quad (13)$$

$$C_{\text{cont}} = 0.0257(10), u = 0.049(15) \quad (N=4). \quad (14)$$

For $N = 6$, there is no evidence of scaling corrections; we find: $C_{\text{cont}} = 0.0236(10)$ for $N = 6$.




FIGURE 2. The scaling ratio $C = \chi/\sigma^2$ versus σ . Dotted lines show the linear fits.

In Fig. 3, our results for C_{cont} are plotted versus $1/N^2$ (the expected order of $1/N$ corrections). They are clearly consistent with a behavior of the type $C_{\text{cont}}(N) = C_{\infty} + B/N^2$. Assuming such a behavior, a fit to $N = 3, 4, 6$ results gives

$$C_{\infty} = 0.0221(13), \quad B = 0.055(19), \quad (15)$$

This is a rather accurate estimate of large- N limit of C , and it is clearly nonzero.

To compare with literature, we set a standard value: $\sqrt{\sigma} = 440$ MeV, for $N \geq 3$. We obtain: $\chi_{\infty}^{1/4} = 170(3)$ MeV, and $\chi^{1/4} = 180(2)$ MeV for $N = 3$. From Eq. (11), with actual values of f_{π} , $m_{\eta'}$, and $N_f = 3$, one has 191 MeV. Using Veneziano's improvement: $m_{\eta'}^2 \rightarrow m_{\eta'}^2 + m_{\eta}^2 - 2m_K^2$, one obtains 180 MeV.

We now turn to b_2 : The cancellations in its definition necessitate very high statistics. We have obtained sufficiently accurate results for $N = 3, 4$, showing scaling. A weighted average, with conservative errors, yields:

$$b_2 = -0.023(7)(N=3), \quad b_2 = -0.013(7)(N=4). \quad (16)$$

Reducing statistical error on b_2 in the $N = 6$ case is extremely difficult. We can only give a rough estimate $b_2 = -0.01(2)$ ($\gamma = 0.348$) in this case, showing that b_2 is very small.




FIGURE 3. The scaling ratio $C = \chi/\sigma^2$ versus $1/N^2$. The line is the linear fit to $A + B/N^2$.

An alternative to cooling is based on the index theorem, using, e.g., the overlap formulation [7]. The rigorous properties of such an approach would make up for a useful consistency check.

In conclusion, we stress that further investigation of other observables in $SU(N)$ theories is called for, to test predictions from: SUSY $SU(N)$, M-theory and gauge/string duality.

REFERENCES

1. B. Lucini and M. Teper, Phys. Lett. B **501**, 128 (2001); Phys. Rev. D **64** (2001) 105019.
2. L. Del Debbio, H. Panagopoulos, P. Rossi, and E. Vicari, Phys. Rev. D **65** (2002) 021501; JHEP **01** (2002) 009; Nucl. Phys. (PS) **106** (2002) 688.
3. L. Del Debbio, H. Panagopoulos, and E. Vicari, JHEP **08** (2002) 044.
4. E. Witten, Ann. Phys. (NY) **128**, 363 (1980).
5. E. Witten, Phys. Rev. Lett. **81**, 2862 (1998).
6. E. Witten, Nucl. Phys. B **156**, 269 (1979); G. Veneziano, Nucl. Phys. B **159**, 213 (1979).
7. H. Neuberger, Phys. Lett. B **417**, 141 (1998).